MARSHALL COUNTY: Key Data Points

- The county observes a significantly lower percentage of unmarried parents compared to the state of Indiana overall, 31.2% versus 37.1% respectively
- The percentage of mothers breastfeeding on discharge is almost 10% higher than the state percentage
- Marshall county observed 25 accidental deaths (14 of which were motor vehicle), 2 suicidal deaths and 1 homicide

Note: All figures relate to 2003 data from the Indiana State Department of Health, unless otherwise specified

Figure 1. Births

	Total		White		Black		Other	
	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*
Marshall County	701	15.1	691	15.2	2	10.5(U)	8	12.1(U)
INDIANA	86,382	13.9	75,422	13.7	9,288	17.5	1,672	10.6

^{*} births per 1,000 population; U = unstable rate, based on fewer than 20 births

Figure 2. Mortality

rigure 2. More	<u>ality</u>							
			NUI	MBER			RATE*	
COUNTY OF RESIDENCE	REPORTED RACE/ ETHNICITY	Live Births	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths (28-364 days)	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths (28-364 days)
MARSHALL	All Races	701	4	4	0	**	**	0
	White	691	4	4	0	**	**	0
	Black	2	0	0	0	0	0	0
	Other	8	0	0	0	0	0	0
	Hispanic	93	0	0	0	0	0	0
INDIANA	All Races	86,382	641	417	224	7.4	4.8	2.6
	White	75,422	479	309	170	6.4	4.1	2.3
	Black	9,288	148	98	50	15.9	10.6	5.4
	Other	1,672	14	10	4	**	**	**
	Hispanic	6,763	38	22	16	5.6	3.3	**

^{*}rate per 1,000 live births **Numerator is less than 20 and the rate is unstable

Figure 3. Outcome Indicators of Mother, by Race


-		V						
		Birth W	/eight			During P		
MARSHALL	Total	% Low	% Very Low	0/ Drotorm	% PNC 1st	0/	%	%
	Total		(% Preterm	_	%	, ,	Unmarried
COUNTY	Births	(< 2500 grams)	grams)	T	Trimester [‡]	Alcohol	Smoked	Parents
All Races	701	5.7	1	7	82.2	1.1	17.4	31.2
White	691	5.6	1	6.9	81.9	1.2	17.4	31.1
Black	2	**	**	**	**	**	**	**

		Birth W	/eight		_	During P		
	Total	% Low	% Very Low	% Preterm	% PNC 1st	%	%	% Unmarried
INDIANA	Births	(< 2500 grams)	(< 1500 grams)	†	Trimester [‡]	Alcohol	Smoked	Parents
All Races	86,382	7.9	1.4	9.9	80.6	0.6	18.5	37.1
White	75,422	7.2	1.2	9.5	82.1	0.6	19.1	32.6
Black	9,288	13.3	3.3	13.3	68.2	1.1	15.2	76.3

[†] Preterm: less than 37 weeks gestation

Figure 4. Breastfeeding

MARSHALL COUNTY	All Races	White	Black
Total Births	701	691	2
Breastfeeding on Discharge	517	511	1
% Breastfeeding on Discharge	73.8	74	50



[‡] PNC: Prenatal Care

'S' Significantly different than the state percent

'**' Outcomes for this county are suppressed due to 20 or fewer total births

Figure 4. Breastfeeding

INDIANA	All Races	White	Black
Total Births	86,382	75,422	9,288
Breastfeeding on Discharge	54,636	49,385	3,959
% Breastfeeding on Discharge	63.2	65.5	42.6




Figure 5. HIV/AIDS

	Total # of AIDS cases reported since 1982*	Total # of HIV cases reported since 1985*
MARSHALL	17	5
COUNTY INDIANA	7,641	3,856

^{*}data gathered from June 2005 HIV/STD Quarterly Report

Figure 6. Nutritional Risk Factors by Status,

Indiana WIC Clinics, Quarter Ending 09/30/2004

Local Agenc	Local Agency: Marshall County Health Department								
Counties Inc	Counties Included: Marshall								
	Childr	Children 1-5 years of Age					%		
Risk Factor or Status	White	Black	Hispanic	Asian	Other	Total	within Agency		
Overweight	14	0	13	0	2	29	3.9%		
At Risk for becoming Overweight	33	0	12	0	3	48	6.4%		

Local Agency: State Wide Counties Included: All

Countres meraded. Am								
	Childr	en 1-5 y		%				
Risk Factor or Status	White	Black	Hispanic	Asian	Other	Total	within Agency	
Overweight	1,732	445	800	12	97	3,086	4.8%	
At Risk for becoming Overweight	2,023	485	639	11	124	3,282	5.1%	

	Priority Issues (Marshall & Indiana)		
Ι	Demographics: Percentage of population enrolled in Medicaid, Percentage of population enrolled in Food Stamps.	Marshall	Indiana
2	% of Population enrolled in Medicaid as of June 2003	10.4	12.4
3	% of Population in Food Stamps Program, 2004	5.4	8.6
P	riority #1 (pregnancy outcomes): Average Infant Mortality Rate, Average Fetal death Rate	Marshall	Indiana
1	Av Total Infant Death Rate (1993-2002)	5.2	8.1
2	Average Fetal Death Rate (1994-2003)	5.4	6.7
	Priority #2 (reduce barriers): Health Professional Shortage Areas, Percentage of women with prenatal care in first trimester	Marshall	Indiana
1	Scores for Primary Care Health Professional Shortage Areas (HPSAs) (2005)	No Score	NA
2	% PNC 1st Trimester†, 2003	82.2	80.6
	Priority #4 (environmental): Prevalence of asthma, Childhood lead poisoning	Marshall	Indiana
1	Prevalence of Asthma in a Medicaid Cohort of Children age 0-17 years, 2003	7.9	10.4
2	%Estimated Lead Blood Levels (%EBLL) for Children under 6 years old FY 2004-2005	NA	2.1
Pı	riority #5 (tobacco): Percentage of mothers who smoked during pregnancy, Percentage of Adult Smoking	Marshall	Indiana
1	% of mothers Smoked During Pregnancy, 2003	17.4	18.5
2	% of Adult Smoking in 2003	29.7	29.3
	Priority #7 (adolescent risk behaviors): Adolescent deaths due to motor vehicle accidents, Chlamydia rates (ages 15-24).	Marshall	Indiana
1	Number of Adolescent (15-24) deaths due to Motor vehicle accidents (2001-2003)	7	
	2003)		748
2	Chlamydia rates per 100,000 county specific population (15-24yrs of Age)	463.43	748 1541.49
2	/		
1	Chlamydia rates per 100,000 county specific population (15-24yrs of Age)	463.43	1541.49
1	Chlamydia rates per 100,000 county specific population (15-24yrs of Age) Priority #8 (obesity): Obesity in pregnant women	463.43 Marshall	1541.49 Indiana
1	Chlamydia rates per 100,000 county specific population (15-24yrs of Age) Priority #8 (obesity): Obesity in pregnant women Number of Obese Pregnant Women In Indiana, 2003 riority #9 (domestic violence, childhood injury): Child deaths due to Injury,	463.43 Marshall 20	1541.49 Indiana 2,496
1 P 1	Chlamydia rates per 100,000 county specific population (15-24yrs of Age) Priority #8 (obesity): Obesity in pregnant women Number of Obese Pregnant Women In Indiana, 2003 riority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates	463.43 Marshall 20 Marshall	1541.49 Indiana 2,496 Indiana
1 Pr 1 2	Chlamydia rates per 100,000 county specific population (15-24yrs of Age) Priority #8 (obesity): Obesity in pregnant women Number of Obese Pregnant Women In Indiana, 2003 riority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates Child deaths due to Injury 1-14yrs (2001-2003)	463.43 Marshall 20 Marshall 3	1541.49 Indiana 2,496 Indiana 257
1 Pr 1 2	Chlamydia rates per 100,000 county specific population (15-24yrs of Age) Priority #8 (obesity): Obesity in pregnant women Number of Obese Pregnant Women In Indiana, 2003 riority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates Child deaths due to Injury 1-14yrs (2001-2003) Child Abuse and Neglect per 1,000 Population under age 18, 2004 Priority #10 (To improve racial and ethnic disparities in women of childbearing age, mothers, and children's health outcomes): Percentage of	463.43 Marshall 20 Marshall 3 16.3	1541.49 Indiana 2,496 Indiana 257 13.5

Source: Data Analysis Team, ISDH. **Notes:** 1. NS= Not Stable 2. NA = Not Available 3. Infant Mortality rate and Fetal Death rate is for 1000 live births. 4. Low Birth Weight: Birth Weight Less than 2,500grams. 5. HPSA Score ranges 1 – 25. The higher the score, the higher the need. 6. EBLL: Ten (10) micrograms of lead per deciliter of whole blood for one venous test; %EBLL is based on screening rate of 27% Statewide, unduplicated kids 7. Obesity in Pregnant Women determined by "Concurrent Illness Obesity" checkbox on the birth certificate 8. PNC percentages are calculated using total births in each county.